arXiv:1206.1170v1 [math.DS] 6 Jun 2012

Reduction of Interaction Delays in Networks

Leonhard Lücken, Jan Philipp Pade, and Serhiy Yanchuk
Humboldt-University of Berlin, Institute of Mathematics,
Unter den Linden 6, 10099, Berlin Germany.

Abstract

Delayed interactions are a common property of coupled natural systems. For instance, signals in neural or laser networks propagate at finite speed giving rise to delayed connections. Such systems are often modeled by delay differential equations with discrete delays. We show that by a selective timeshift transformation it is often possible to reduce the number of different delays. We identify dynamic invariants of this transformation, determine its capabilities to reduce the number of delays and interpret these findings in terms of the topology of the underlying graph. In particular, we show that networks with identical generalized round-trip times along the fundamental semicycles are dynamically equivalent.

Time delays play an important role in networks of coupled dynamical systems. For instance, in neuronal networks [1–5] delays occur due to the finite propagation times along the axons or to reaction times at chemical synapses. Similarly, the propagation of light between interacting lasers [6–8] may cause significant time delays. Depending on the physiological properties of axons and synapses, the delays of signals between different cells in the network differ and in practice there are as many different delays in the network as there are links between single neurons. Such a diversity of delays was shown to enable the design of robust pattern generating systems [9, 10] and to play an important role in neural processing of temporal information [11]. Hence, to study such natural systems, one would need to study models involving numerous different delays. Independently of the physical origin, dynamics in networks of interacting systems can often be described by the equations

$$\dot{x}_j(t) = f_j\left(x_j(t), \left(x_k\left(t-\tau_{jk}\right)\right)_{k\in P_j}\right), \tag{1}$$

where $x_j(t)$, $j = 1, \ldots, N$, denotes the dynamics of a node j and P_j is the set of all other nodes of the network connected to the node j. For instance, x_j may be considered as a dynamical variable describing the dynamics of one neuron in the network. Taking into account finite time delays τ_{jk}, which occur due to the signal propagation time from a node k to the node j, the coupling terms contain all variables x_k which are connected to x_j with corresponding time delays τ_{jk}. Here we consider directed networks, thus one may generally have $\tau_{jk} \neq \tau_{kj}$ or even $k \in P_j$ while $j \notin P_k$. We assume that the network is connected and hence, the number L of links is larger than $N - 1$.

For a network of N systems, up to N^2 different time delays may occur in (1). By allowing several connections from one system to another, the number of delays can increase even more. This creates immense challenges for the analysis of the system, since every single delay may alter the properties and dynamics significantly [12–16]. The interaction of several delays is even more complicated [8–10, 17, 18] and not fully understood so far.

In this Letter we show how the number of different delays can be reduced. It appears that any connected network possesses a characteristic number of delays which is essential for describing the dynamics. This number of essential delays equals the cycle space dimension $C = L - (N - 1)$ of the underlying graph and is usually smaller than the number of distinct τ_{jk}. We show that the essential delays correspond to generalized round-trip times along fundamental semicycles in the network. Networks which have the same local dynamics and the same set of essential delays can be considered as equivalent from the dynamical point of view.

As a simple illustrative example, let us consider a ring of unidirectionally coupled systems

$$\dot{x}_j(t) = f\left(x_j(t), x_{j+1}(t-\tau_j)\right) \tag{2}$$

with inhomogeneous coupling delays τ_j. It is known [19, 20] that this system can be reduced to the ring where all time delays are equal to $\tau = \frac{1}{N}\sum\tau_j$. This can be done by a separate time-shift transformation for each node

$$y_j(t) = x_j(t+\eta_j), \quad 1 \leq j \leq N. \tag{3}$$

Indeed, the transformed variables $y_j(t)$ fulfill

$$\dot{y}_j(t) = f\left(y_j(t), y_{j+1}(t-\tau_j+\eta_j-\eta_{j+1})\right),$$

and if the shifts η_j are chosen appropriately, all hitherto distinct delays become the same while their sum along the ring, the round-trip time $N\tau$, is preserved. This homogenization of delays was applied to simplify the analysis of the solutions in systems (2) with inhomogeneous delays [20, 21]. Conversely, the same transformation was applied to design pattern generators [9, 10] starting from the system with homogeneous delays. Our aim is to use the time-shift transformation (3) to reduce the number of different delays in networks with arbitrary topology.

Applying (3) to (1), we obtain the new system

$$\dot{y}_j(t) = f_j\left(y_j(t), \left(y_k(t-\tilde{\tau}_{jk})\right)_{k\in P_j}\right) \tag{4}$$

with $\tilde{\tau}_{jk} = \tau_{jk} + \eta_k - \eta_j$. We proceed by exploring possible delay reductions in basic network motifs, which consist of exactly one semicycle (a cycle in the underlying undirected graph) and are depicted in Fig. 1.

Motif I. Dynamics on network I is described by the system of equations

$$\begin{aligned}\dot{x}_1(t) &= f_1\left(x_1(t)\right),\\ \dot{x}_2(t) &= f_2\left(x_2(t), x_1(t-\tau_1), x_1(t-\tau_2)\right).\end{aligned}$$

The transformation (3) leads to the same system for y_j but with different delays $\tilde{\tau}_j = \tau_j + \eta_1 - \eta_2,\ j = 1, 2$. Without loss of generality we assume $\tau_1 < \tau_2$. By choosing $\eta_2 - \eta_1 = \tau_1$,



Figure 1: Simple motifs consisting of one semicycle.

we obtain $\tilde{\tau}_1 = 0$ and $\tilde{\tau}_2 = \tau_2 - \tau_1 > 0$. Thus, the transformed system contains only one delayed connection – the other one becomes instantaneous. In case $\tau_1 = \tau_2$ we obtain $\tilde{\tau}_1 = \tilde{\tau}_2 = 0$, which means that both delays can be eliminated. Thus, the dynamics in motif I for $\tau_1 \neq \tau_2$ can be described by a system with just one delay, and for $\tau_1 = \tau_2$ the system reduces to an ordinary differential equation. Note that the quantity $\tilde{\tau}_2 - \tilde{\tau}_1 = \tau_2 - \tau_1$ remains invariant.

Motif II. After applying transformation (3) to motif II, the delays change to $\tilde{\tau}_1 = \tau_1 + \eta_1 - \eta_2$, $\tilde{\tau}_2 = \tau_2 + \eta_1 - \eta_3$, $\tilde{\tau}_3 = \tau_3 + \eta_2 - \eta_3$. It is easy to check that by appropriate choice of η_j one obtains a system of equations with a single delay. For instance, it can be done by requiring that all new delays are identical $\tilde{\tau}_1 = \tilde{\tau}_2 = \tilde{\tau}_3 \equiv \tau$. In this case the delay time is $\tau = \tau_1 - \tau_2 + \tau_3$. The corresponding time shifts can be chosen as $\eta_1 = \mu$, $\eta_2 = \mu + \tau_2 - \tau_3$, $\eta_3 = \mu + 2\tau_2 - \tau_1 - \tau_3$ with an arbitrary constant μ. Simple calculations give five different reductions to a single delay as shown in Table I. Note that some of the reductions lead to negative delays in the resulting network, which should be avoided since the dynamics in such systems are unstable [22]. For example, in the case $\tau = \tau_1 - \tau_2 + \tau_3 < 0$ there is only one suitable reduction ($\tilde{\tau}_1 = \tilde{\tau}_3 = 0$ and $\tilde{\tau}_2 = -\tau$). Summarizing, the motif II always admits a reduction to one delay $T = |\tau_1 - \tau_2 + \tau_3| = |\tilde{\tau}_1 - \tilde{\tau}_2 + \tilde{\tau}_3|$. This invariant quantity can be considered as a generalized round-trip time, taking into account the directionality of the connections. In the case $\tau = 0$ the delays can be eliminated.

Motif III. Similarly, for the motif III [Fig. 1-III], one can reduce the four different delays τ_1,τ_2, τ_3, and τ_4 to a single delay τ. Formally, ten different combinations are possible, which are listed in Table I(c). As in the case of motif I a reduction to uniform delays at each link is impossible for this case. But, as in the other examples, a reduction to a system with only one delayed connection can be achieved. Further, the generalized round-trip $T = |\tau_1 - \tau_2 - \tau_3 + \tau_4|$ is preserved for any reduction. When this round-trip is zero, all (a): motif I

$\tilde{\tau}_1$	0	τ
$\tilde{\tau}_2$	$-\tau$	0

(b): motif II

$\tilde{\tau}_1$	0	0	τ	$\tau/2$	τ
$\tilde{\tau}_2$	0	$-\tau$	0	0	τ
$\tilde{\tau}_3$	τ	0	0	$\tau/2$	τ

(c): motif III

$\tilde{\tau}_1$	0	0	0	0	0	τ	$\tau/2$	τ	τ	$-\tau$
$\tilde{\tau}_2$	0	0	$-\tau$	$-\tau/2$	$-\tau$	0	0	0	τ	$-\tau$
$\tilde{\tau}_3$	0	$-\tau$	0	$-\tau/2$	$-\tau$	0	0	τ	0	$-\tau$
$\tilde{\tau}_4$	τ	0	0	0	$-\tau$	0	$\tau/2$	τ	τ	0

Table I: Possible reductions to a single delay in motifs I, II, and III [see Fig. 1]. The values of the delay τ are as follows: $\tau = \tau_1 - \tau_2$ (a), $\tau = \tau_1 - \tau_2 + \tau_3$ (b), $\tau = \tau_1 - \tau_2 - \tau_3 + \tau_4$ (c). In each case, the generalized round-trip is given by $T = |\tau|$.

delays can be eliminated.

General networks. The cases considered above suggest that the generalized round-trip time along a semicycle in the network remains invariant under the time shift transformation (3). Indeed, this holds for arbitrary networks. Let us first define a semicycle $c = (\ell_1, ..., \ell_k)$ of a directed graph as a closed undirected path, which is represented as an ordered set of its constituting links. Then we define the generalized round-trip time $T(c)$ along a given semicycle $c = (\ell_1, ..., \ell_k)$ as follows

$$T(c) := \left| \sum_{j=1}^{k} \Gamma_j \tau(\ell_j) \right| . \tag{5}$$

Here $\tau(\ell_j)$ is the time delay along the link ℓ_j and Γ_j is either $+1$ or -1 depending on the direction of the link ℓ_j with respect to an arbitrary but fixed orientation of the semicycle. Note that the orientation can be chosen in two different ways, which corresponds to opposite signs of Γ_j. The obtained value for $T(c)$ is independent of the choice of orientation.

It is possible to prove that there is a choice of timeshifts η_j and a set $S = \{\ell_1, ..., \ell_{N-1}\}$ of $N-1$ links, such that the delays $\tilde{\tau}_{jk} = \tau_{jk} - \eta_j + \eta_k$ in the transformed equation (4) are zero on all links contained in S and non-negative on all other links. In fact, the set S is a spanning tree: a set of $N-1$ links which does not contain any semicycle. The addition of any link $\ell \notin S$ results in a set $S' = S \cup \{\ell\}$ which contains exactly one semicycle. This is the fundamental semicycle corresponding to ℓ. The transformed delay $\tilde{\tau}(\ell)$ on the link ℓ



Figure 2: (a) A network with cycle space dimension $C = 5$. A spanning tree S [solid links] and one of its fundamental cycles, $c(\ell)$ [blue dotted path], are displayed. The ±-signs indicate the value of the corresponding coefficients Γ_j in the the round-trip $T(c(\ell))$, see (5). (b) A network of $N = 6$ cells with cycle space dimension $C = 3$. We depict a spanning tree [solid links] and three corresponding fundamental cycles which are indicated by differently dashed paths. The numbers refer to an indexing of the links which is used in the text. According to our results, any admissible dynamics in this network involves at most three different interaction delays.

equals the round-trip delay of this fundamental semicycle [Fig. 2(a)]. To prove this claim one can inductively construct timeshifts and the corresponding spanning tree, which contains instantaneous links. Details will be presented elsewhere [23]. Consequently, the number of delays in the transformed system can be reduced to at most

$$C = L - (N - 1). \tag{6}$$

The number C has a well known meaning for network graphs [24]: It is the cycle space dimension which is defined as the maximal number of independent cycles. Here, a set of cycles is called independent if neither of the cycles is obtained as a sequence of the other ones in the set. For instance, in the network given in Fig. 2(a) the cycle space dimension is $C = 5$ and in the one depicted in (b) it is $C = 3$.

We call C the *essential number* of delays since it is the minimal number of different delays to which the number of delays in a network can be reduced generically. Here, generically means that the conditions which allow for further reduction of delays form a nullset in the parameter space of delays $\mathbb{R}^L_{\geq 0} = \{\tau_{jk} \mid \tau_{jk} \geq 0\}$ of the original system (1) [23].

Dynamical invariants. There is a natural correspondence between solutions of the original system (1) and the transformed system (4): Any solution $\mathbf{x}(t) = (x_j(t))_{1 \leq j \leq N}$ of (1) can



Figure 3: Equivalent stable solutions of unidirectional rings of four Mackey-Glass oscillators (7) with distinct single delays but equal round-trip. In each row, the time evolution of the j-th cell is indicated by the coloring. In (a), the delays are homogeneous, $\tau_{j,j+1} = 25$. Chart (b) shows the corresponding solution of the transformed system with delays $\tilde{\tau}_{j,j+1} = \tau_{j,j+1} - \eta_j + \eta_{j+1}$ and $\eta = (-5.8, 0, -5.6, -18.4)$. Other parameters: $\beta = 0.2$, $\gamma = 0.1$, $c = 4$.

be identified with its transformation, $\mathbf{y}(t) = (x_j(t + \eta_j))_{1 \le j \le N}$, which is a solution of (4), see Fig. 3. As a result, many important properties of solutions like stability, frequency of oscillation, amplitude, etc., remain the same within a class of equivalent systems. They depend only on a set of generalized round-trip time delays, but not on the specific delays τ_{jk}. As a consequence it is possible (and favorable) to perform bifurcation analysis in the reduced parameter space. Practically, one considers round-trips on a set of fundamental semicycles instead of single delays as parameters of the system.

The following example is intended to provide an idea of possible applications and merits of the results presented in this article. Let us consider the network of Mackey-Glass systems [7, 25, 26]

$$\dot{x}_j(t) = -\gamma x_j(t) + \beta \frac{c \sum x_k(t - \tau_{jk})}{1 + (c \sum x_k(t - \tau_{jk}))^{10}}, \tag{7}$$

where the summation is performed over all coupled nodes $k \in P_j$. The coupling topology is shown in Fig. 2(b). The parameters are fixed to $\gamma = 0.1$, $c = 0.525$ and $\beta = 0.2$. Although the total number of delays is 8, one can parametrize all of them by the three round-trips $T(c_j)$, $j = 1, 2, 3$, on the fundamental semicycles [Fig. 2(b)]. For further investigation we fix $T(c_3) = 7$ and $T(c_2) = 10$ wherever only $T(c_1)$ is varied.

Firstly, the delays in the network were set up randomly in such a way that the round-trips were preserved with $T(c_1) = 5$. For each realization of delays, the system was integrated numerically with constant initial conditions [see caption to Fig. 4]. The empirical distribution of the individual delays and the observed attractors is shown in Fig. 4(b). One can observe two attractors, one equilibrium and one periodic solution, which exist independently of the choice of single delays. They can be identified by their componentwise supremum norm $|x| = \sqrt{\sum_{j=1}^{N} \sup_t |x_j(t)|^2}$ which is invariant under (3). Starting from those solutions, we obtain the bifurcation diagram with respect to $T(c_1)$ [Fig. 4(a)]. For $T(c_1) = 0$, two stable equilibria are the only attractors for the system. The lower equilibrium looses stability in a Hopf bifurcation (H) and the emergent periodic solution undergoes a period doubling bifurcation (P1) which connects to a reverse period doubling at (P2). Snaking of the main branch accompanied by period doubling bridges and divergent periods indicates the presence of a homoclinic saddle-focus [27]. A step further can be done by a two-parametric bifurcation diagram [Fig. 4(c)]. There, the Hopf bifurcation (H) was followed numerically in the region $0 \leq T(c_1) \leq 500$ and $0 \leq T(c_2) \leq 500$. Allthough the same diagram might have been obtained accidentally by studying variations of the delays τ_5 and τ_7, its proper meaning as a diagram of codimension one is first made accessible by the presented results.

Discussion. We have presented a universal method for the reduction of discrete interaction-delays in networks of dynamical systems. It turned out that the cycle space dimension of the network equals the number of essential delays which determine the dynamics. The networks whose generalized round-trip times along the fundamental semicycles coincide, form a class of dynamically equivalent systems. This is an important step towards the systematization and classification of dynamics on networks with many different delays. Note that all results which are presented in this paper can be extended straightforwardly to networks with multiple links from one cell to another (see, for instance, motif I in Fig. 1) and to other types of local dynamics like time-discrete systems or evolution equations.

We thank K. Knauer and M. Zaks for useful discussions and the DFG for financial support in the framework of the Collaborative Research Center SFB910.

---

[1] C. Leibold and J. L. van Hemmen, Phys. Rev. Lett. **94**, 168102 (2005).

[2] E. Rossoni, Y. Chen, M. Ding, and J. Feng, Phys. Rev. E **71**, 061904 (2005).

[3] R. M. Memmesheimer and M. Timme, Physica D **224**, 182 (2006).



Figure 4: Dynamical features of six-node network in Fig. 2(b) of Mackey-Glass systems (7). (a): Bifurcation diagram for $T(c_1) \in [0, 23]$ and $T(c_2) = 10$. Black lines correspond to stable solutions and gray to unstable. Solid branches are equilibria, dashed are periodic solutions. Bifurcations are denoted by 'H' (Hopf), 'P[n]' (period doubling), and 'F[n]' (fold). The ordinate measures the norm $|x|$ and the inset shows the period along the main branch. Red crosses indicate the observed solutions from chart (b). (b): 50 random realizations of single delays which keep constant $T(c_1) = 5$, $T(c_2) = 10$, and $T(c_3) = 7$. Each dot corresponds to a realization of τ_j. Mean and standard deviation are indicated by squares and bars. For each choice of delays, constant initial functions were drawn from a uniform distribution on $[0, 1.5]$. The inset bars indicate the number of observed solutions $x(t)$, $t \in [10^5, 1.1 \times 10^5]$, with the norm $|x|$ in their base interval. Chart (c) shows the trace and reappearances of the Hopf bifurcations in the $T(c_1)$-$T(c_2)$-plane.

[4] T.-W. Ko and G. B. Ermentrout, Phys. Rev. E **76**, 056206 (2007).

[5] D. J. Bakkum, Z. C. Chao, and S. M. Potter, PLoS ONE **3**, e2088 (2008).

[6] R. Vicente, S. Tang, J. Mulet, C. R. Mirasso, and J.-M. Liu, Phys. Rev. E **73**, 047201 (2006).

[7] O. D'Huys, R. Vicente, T. Erneux, J. Danckaert, and I. Fischer, Chaos **18**, 037116 (2008).

[8] A. L. Franz, R. Roy, L. B. Shaw, and I. B. Schwartz, Phys. Rev. E **78**, 016208 (2008).

[9] S. Yanchuk, P. Perlikowski, O. V. Popovych, and P. A. Tass, Chaos **21**, 047511 (2011).

[10] O. V. Popovych, S. Yanchuk, and P. A. Tass, Phys. Rev. Lett. **107**, 228102 (2011).

[11] C. E. Carr, Annu. Rev. Neurosci. **16**, 223 (1993).

[12] T. Erneux, *Applied Delay Differential Equations*, vol. 3 (Springer, 2009).

[13] J. Foss, A. Longtin, B. Mensour, and J. Milton, Phys. Rev. Lett. **76**, 708 (1996).

[14] G. Giacomelli and A. Politi, Phys. Rev. Lett. **76**, 2686 (1996).

[15] V. Flunkert, S. Yanchuk, T. Dahms, and E. Schöll, Phys. Rev. Lett. **105**, 254101 (2010).

[16] S. Heiligenthal, T. Dahms, S. Yanchuk, T. Jüngling, V. Flunkert, I. Kanter, E. Schöll, and W. Kinzel, Phys. Rev. Lett. **107**, 234102 (2011).

[17] J. K. Hale and S. M. Tanaka, J. Dyn. Differ. Equ. **12**, 1 (2000).

[18] I. Kanter, M. Zigzag, A. Englert, F. Geissler, and W. Kinzel, Europhys. Lett. **93**, 60003 (2011).

[19] G. V. der Sande, M. C. Soriano, I. Fischer, and C. R. Mirasso, Phys. Rev. E **77**, 055202 (pages 4) (2008).

[20] P. Perlikowski, S. Yanchuk, O. V. Popovych, and P. A. Tass, Phys. Rev. E **82**, 036208 (2010).

[21] P. Baldi and A. Atia, IEEE Transactions on Neural Networks **5**, 1045 (1994).

[22] J. K. Hale and S. M. V. Lunel, *Introduction to Functional Differential Equations* (Springer-Verlag, 1993).

[23] L. Lücken, J. Pade, K. Knauer, and S. Yanchuk, in preparation (2012).

[24] R. Diestel, *Graph Theory* (Springer, 2010).

[25] M. C. Mackey and L. Glass, Science **197**, 287 (1977).

[26] L. Appeltant, M. C. Soriano, G. Van der Sande, J. Danckaert, S. Massar, J. Dambre, B. Schrauwen, C. R. Mirasso, and I. Fischer, Nat. Commun. **2** (2011).

[27] P. Glendinning and C. Sparrow, J. Stat. Phys. **35**, 645 (1984).